

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 5: C12N 5/00, 15/00, A01H 1/04, 5/00, 5/10		A1	(11) International Publication Number: WO 94/17176
		(43) International Publication Date: 4 August 1994 (04.08.94)	
(21) International Application Number: PCT/US94/00927		(81) Designated States: JP, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).	
(22) International Filing Date: 27 January 1994 (27.01.94)		Published With international search report. With amended claims and statement.	
(30) Priority Data: 08/010,997 29 January 1993 (29.01.93) US			
(71) Applicant: PURDUE RESEARCH FOUNDATION [US/US]; Division of Sponsored Programs, Room 328, 1650 Engineering Administration Building, West Lafayette, IN 47907-1650 (US).			
(72) Inventors: HODGES, Thomas, K.; 601 Waldron Street, West Lafayette, IN 47906 (US). LYZNIK, Leszek, A.; 2550 Yeager Road 10-8, West Lafayette, IN 47906 (US).			
(74) Agents: LAMMERT, Steven, R. et al.; Barnes & Thornburg, 1313 Merchants Bank Building, 11 South Meridian Street, Indianapolis, IN 46204 (US).			

(54) Title: CONTROLLED MODIFICATION OF EUKARYOTIC GENOMES

(57) Abstract

DNA constructs are provided for the creation of transgenic eukaryotic cells. The figure provides a schematic representation of the DNA constructs used in accordance with this invention. These DNA constructs allow a more precise and effective transformation procedure by enabling the targeting of DNA sequences for insertion into a particular DNA locus, while enabling the removal of any randomly inserted DNA sequences that occur as a by-product of known transformation procedures.



- ▷ 1st RECOMBINATION TARGET SEQUENCE
- ▨ HOMOLOGOUS REGION
- ▭ 1st SELECTABLE MARKER GENE
- ▤ DNA SEQUENCE OF INTEREST
- 2nd RECOMBINATION TARGET SEQUENCE
- △ POLYLINKER REGION
- ▽ 2nd RECOMBINASE GENE
- ▽ 1st RECOMBINASE GENE
- a - DNA TARGETED FOR INSERTION VIA HOMOLOGOUS RECOMBINATION
- b - MULTIFUNCTIONAL DNA SEQUENCE
- c - EXCISABLE SELECTION REGION

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

-1-

CONTROLLED MODIFICATION OF EUKARYOTIC GENOMESBackground and Summary of the Invention

5 This invention relates to a method of transforming eukaryotic cells. More particularly, this invention relates to the use of DNA constructs designed to insert a particular DNA fragment into a chosen locus of the host cell's DNA.

10 Through the use of recombinant DNA technology, foreign DNA sequences can be inserted into an organism's genome to alter the phenotype of the organism. A variety of different procedures have been described and utilized to produce stably transformed eukaryotic cells. All of these procedures are based on first introducing the foreign DNA
15 into the eukaryotic cell, followed by isolation of those cells containing the foreign DNA integrated into the eukaryotic cell's DNA.

Unfortunately, all current higher eukaryotic cell transformation procedures produce transformed cells that
20 contain the introduced foreign DNA inserted randomly and throughout the genome and in multiple copies. This random insertion of introduced DNA into the genome of host cells can be lethal if the foreign DNA happens to insert into, and thus mutate, a critically important native gene. In
25 addition, even if the random insertion event does not impair the functioning of a host cell gene, the expression of an inserted foreign gene may be influenced by "position effects" caused by the surrounding genomic DNA. Thus a gene could be inserted into a site where the position
30 effects are strong enough to prevent the synthesis of an effective amount of product from the introduced gene. Finally, because over-production of a gene product can have deleterious effects on the cell, genetic engineers typically wish to control the copy number of the inserted

-2-

DNA, often only desiring the insertion of a single copy of the DNA sequence.

For these reasons a transformation system is desired that allows a gene to be targeted to a specific site of the host's genome. Preferably, such a system would also provide a means of preventing or subsequently removing any randomly inserted DNA sequences.

The present invention enables the targeting of a length of DNA to a specific non-lethal site in the host cell's genome, and provides for the removal of any randomly inserted DNA sequences. In addition, a proper selection of the targeted site can minimize position effects, enabling an inserted gene to synthesize an effective amount of its protein product. Therefore, this invention allows a much more precise and effective system of genetic engineering than is currently possible.

Additional objects, features, and advantages of the invention will become apparent to those skilled in the art upon consideration of the following detailed description of preferred embodiments exemplifying the best mode of the invention as presently perceived.

Brief Description of the Drawings

Figs. 1A and 1B are schematic representations of DNA constructs in accordance with this invention.

Figs. 2A and 2B illustrate plasmid constructions and general strategies for recombination experiments.

Figs. 3A and 3B are diagrammatic representations of FLP expression vectors used to study activity of yeast FLP/FRT system in plant protoplasts.

Figs. 4A and 4B illustrate the structure of the FRT site-containing vectors.

Fig. 5 depicts vectors useful for testing integrative recombination.

- 3 -

Fig. 6 provides diagrams of vectors for testing excisional recombination.

Fig. 7 illustrates the general strategy for a positive selection of gene targeting events using the FLP/FRT site-specific recombination system.

Fig. 8 provides a schematic map for a targeting vector in accordance with the present invention.

Fig. 9 illustrates examples of DNA constructs in accordance with this invention.

10

Detailed Description of the Invention

In accordance with one preferred embodiment of the present invention, eukaryotic cells are transformed with DNA sequences that are targeted to a predetermined sequence of the eukaryote's DNA. Typically the introduced DNA sequences will constitute entire functional genes.

Eukaryotic cells can also be transformed with other DNA sequences such as gene transcription and translation regulatory sequences. Transcription and translation regulatory sequences are those DNA sequences necessary for efficient expression of a gene product. In general such regulatory elements can be operably linked to any gene to control the gene's expression, the entire unit being referred to as the "expression cassette." An expression cassette will typically contain, in addition to the coding sequence, a promoter region, a translation initiation site and a translation termination sequence. Unique endonuclease restriction sites may also be included at the ends of an expression cassette to allow the cassette to be easily inserted or removed when creating DNA constructs.

The expression of a gene is primarily directed by its own promoter, although other DNA regulatory elements are necessary for efficient expression of a gene product. Promoter sequence elements include the TATA box consensus

-4-

sequence (TATAAT), which is usually 20 to 30 base pairs (bp) upstream of the transcription start site. In most instances the TATA box is required for accurate transcription initiation. By convention, the transcription start site is designated +1. Sequences extending in the 5' (upstream) direction are given negative numbers and sequences extending in the 3' (downstream) direction are given positive numbers.

Promoters can be either constitutive or inducible. A constitutive promoter controls transcription of a gene at a constant rate during the life of a cell, whereas an inducible promoter's activity fluctuates as determined by the presence (or absence) of a specific inducer. The regulatory elements of an inducible promoter are usually located further upstream of the transcriptional start site than the TATA box. Ideally, for experimental purposes, an inducible promoter should possess each of the following properties: a low to nonexistent basal level of expression in the absence of inducer, a high level of expression in the presence of inducer, and an induction scheme that does not otherwise alter the physiology of the cell. The basal transcriptional activity of all promoters can be increased by the presence of "enhancer" sequences. Although the mechanism is unclear, certain defined enhancer regulatory sequences are known, to those familiar with the art, to increase a promoter's transcription rate when the sequence is brought in proximity to the promoter.

The creation of a transformed cell requires that the DNA be physically placed within the host cell. Current transformation procedures utilize a variety of techniques to introduce DNA into a cell. In one form of transformation, the DNA is microinjected directly into cells through the use of micropipettes. Alternatively, high velocity ballistics can be used to propel small DNA associated particles into the cell. In another form, the

-5-

cell is permeabilized by the presence of polyethylene glycol, thus allowing DNA to enter the cell through diffusion. DNA can also be introduced into a cell by fusing protoplasts with other entities which contain DNA.

- 5 These entities include minicells, cells, lysosomes or other fusible lipid-surfaced bodies. Electroporation is also an accepted method for introducing DNA into a cell. In this technique, cells are subject to electrical impulses of high field strength which reversibly permeabilizes biomembranes, allowing the entry of exogenous DNA sequences.
- 10

- In addition to these "direct" transformation techniques, transformation can be performed via bacterial infection using *Agrobacterium tumefaciens* or *Agrobacterium rhizogenes*. These bacterial strains contain a plasmid (called Ti or Ri respectively) which is transmitted into plant cells after infection by *Agrobacterium*. One portion of the plasmid, named transferred DNA (T-DNA), is then integrated into the genomic DNA of the plant cell. This system has been extensively described in the literature and can be modified to introduce foreign genes and other DNA sequences into plant cells.
- 15
- 20

- Agrobacterium*-mediated transformation works best with dicotyledonous diploid plant cells whereas the direct transformation techniques work with virtually any cell.
- 25 Direct transformation techniques can be used to transform haploid cells obtained from immature inflorescences of plants.

- Transformed cells (those containing the DNA inserted into the host cell's DNA) can be selected from untransformed cells if a selectable marker was included as part of the introduced DNA sequences. Selectable markers include genes that provide antibiotic resistance or herbicide resistance. Cells containing these genes are capable of surviving in the presence of antibiotic or herbicide concentrations that kill untransformed cells.
- 30
- 35

-6-

Examples of selectable markers include the *bar* gene which provides resistance to the herbicide Basta, the *nptII* gene which confers kanamycin resistance and the *hpt* gene which confers hygromycin resistance.

5 Once a transformed plant cell is generated, an entire plant can be obtained through cell culturing techniques. Individual cultured cells divide to give rise to an undifferentiated mass of cells called callus tissue. Once callus tissue is formed, shoots and roots may be
10 induced from the callus by techniques known to those familiar with the art, and the resulting plantlets can be planted. Alternatively, somatic embryo formation can be induced in the callus tissue. These somatic embryos germinate as natural embryos to form plants. All the
15 transformation techniques described above have the limitation that they result in multiple copies inserted, and (but to a lesser extent in *Agrobacterium*-mediated transformations) result in foreign DNA insertions throughout the genome.

20 The present invention enables the targeting of a length of DNA to a specific site in the host cell's genome that is non-lethal, and provides for the removal of any introduced DNA sequences randomly inserted into the genome. Presumably the selected target site will also allow the
25 inserted gene to produce its protein product in an amount sufficient to produce the desired effect.

 The first element of the invention involves targeting the gene to a specific site in the host cell's genome. Targeting will be carried out via "homologous
30 recombination." Homologous recombination is a reaction between any pair of DNA sequences having a similar sequence of nucleotides (homologous sequences), where the two sequences interact (recombine) to form a new recombinant DNA species. The frequency of homologous recombination
35 increases as the the length of the shared nucleotide DNA

-7-

sequences increases, and is higher with linearized plasmid molecules than with circularized plasmid molecules.

Homologous recombination can occur between two DNA sequences that are less than identical, but the

- 5 recombination frequency declines as the divergence between the two sequences increases.

- Introduced DNA sequences can be targeted via homologous recombination by linking the DNA of interest to sequences sharing homology with endogenous sequences of the
- 10 host cell. Once the DNA enters the cell, the two homologous sequences can interact to insert the introduced DNA at the site where the homologous genomic DNA sequences were located. Therefore, the choice of homologous sequences contained on the introduced DNA, will determine
- 15 the site where the introduced DNA is integrated via homologous recombination. For example, if the DNA sequence of interest is linked to DNA sequences sharing homology to a single copy gene of the host eukaryotic cell, the DNA sequence of interest will be inserted via homologous
- 20 recombination at only that single specific site. However, if the DNA sequence of interest is linked to DNA sequences sharing homology to a multi-copy gene of the host eukaryotic cell, then the DNA sequence of interest can be inserted via homologous recombination at each of the
- 25 specific sites where a copy of the gene is located. Preferable, the predetermined host DNA site of insertion is a non-essential endogenous gene present in high copy number, e.g. a storage protein gene.

- DNA can be inserted into the genome by a
- 30 homologous recombination reaction involving either a single reciprocal recombination (resulting in the insertion of the entire length of the introduced DNA) or through a double reciprocal recombination (resulting in the insertion of only the DNA located between the two recombination events).
- 35 For example if one wishes to insert a foreign gene into the

- 8 -

genomic site where the gene encoding alcohol dehydrogenase (the *adh* gene) is located, the introduced DNA should contain sequences homologous to the genomic *adh* gene. A single homologous recombination event would then result in the entire introduced DNA sequence being inserted into the genomic *adh* gene. Alternatively, a double recombination event can be achieved by flanking each end of the DNA sequence of interest (the sequence intended to be inserted into the genome) with DNA sequences homologous to the *adh* gene. A homologous recombination event involving each of the homologous flanking regions will result in the insertion of the foreign DNA. Thus only those DNA sequences located between the two regions sharing genomic homology become integrated into the genome.

Although introduced sequences can be targeted for insertion into a specific genomic site via homologous recombination, in higher eukaryotes homologous recombination is a relatively rare event compared to random insertion events. In plant cells, foreign DNA molecules find homologous sequences in the cell's genome and recombine at a frequency of $0.5 - 4.2 \times 10^{-4}$ (the number of targeted events divided by the number of random integrations events). Thus any transformed cell that contains an introduced DNA sequence integrated via homologous recombination will also likely contain numerous copies of randomly integrated introduced DNA sequences. Therefore to maintain control over the copy number and the location of the inserted DNA, these randomly inserted DNA sequences must be removed.

One manner of removing these random insertions is to utilize a site-specific recombinase system. In general, a site specific recombinase system consist of three elements: two pairs of DNA sequence (the site-specific recombination sequences) and a specific enzyme (the site-specific recombinase). The site-specific recombinase will

-9-

catalyze a recombination reaction only between two site-specific recombination sequences.

Depending on the orientation of the site-specific recombination sequences, intervening sequences will either be excised or inverted in the presence of the site specific recombinase. When the site-specific recombination sequences are orientated in opposite directions relative to one another (ie, inverted repeats) then any intervening sequences will be inverted relative to the other sequences in the genome. However, if the site-specific recombination sequences are orientated in the same direction relative to one another (ie, direct repeats) any intervening sequences will be deleted upon interaction with the site specific recombinase. Therefore, if direct repeats of the site-specific recombination sequences are present at both ends of DNA introduced into the cell, any random integration of these sequences can be subsequently removed by interaction of the site-specific recombination sequences with their site specific recombinase. Unlike the random integration event, a homologous recombination event will only integrate a portion of the introduced DNA. If this portion of the DNA does not contain both site-specific recombination sequences, then even in the presence of the site specific recombinase these sequences will remain integrated while the randomly integrated DNA sequences will be excised.

A number of different site specific recombinase systems can be used, including but not limited to the Cre/lox system of bacteriophage P1, the FLP/FRT system of yeast, the Gin recombinase of phage Mu, the Pin recombinase of E. coli, and the R/RS system of the pSR1 plasmid. The two preferred site specific recombinase systems are the bacteriophage P1 Cre/lox and the yeast FLP/FRT systems. In these systems a recombinase (Cre or FLP) will interact specifically with its respective site-specific recombination sequence (lox or FRT respectively) to invert

-10-

or excise the intervening sequences. The sequence for each of these two systems is relatively short (34 bp for lox and 47 bp for FRT). Currently the FLP/FRT system of yeast is the preferred site specific recombinase system since it normally functions in a eukaryotic organism (yeast), and is well characterized. Applicants have reason to believe that the eukaryotic origin of the FLP/FRT system allows the FLP/FRT system to function more efficiently in eukaryotic cells than the prokaryotic site specific recombinase systems.

The FLP/FRT recombinase system has been demonstrated to function efficiently in plant cells. Experiments on the performance of the FLP/FRT system in both maize and rice protoplasts indicates that FRT site structure, and amount of the FLP protein present, affects excision activity. In general, short incomplete FRT sites leads to higher accumulation of excision products than the complete full-length FRT sites. Site specific recombination systems can catalyze both intra- and intermolecular reactions in maize protoplasts, indicating that the system can be used for DNA excision as well as integration reactions. The recombination reaction is reversible and this reversibility can compromise the efficiency of the reaction in each direction. Altering the structure of the site-specific recombination sequences is one approach to remedying this situation. The site-specific recombination sequence can be mutated in a manner that the product of the recombination reaction is no longer recognized as a substrate for the reverse reaction, thereby stabilizing the integration or excision event.

Another approach to manipulate the system is based on the entropic advantage of a unimolecular (excision) over a bimolecular (integration) reaction. By limiting the expression of the recombinase enzyme, the efficiency of the integrative recombination, the

-11-

thermodynamically least favored event, can be reduced. Experiments in maize protoplasts indicate higher concentration of the FLP protein increased the efficiency of the excision reaction.

5 In one embodiment, a strong monocot promoter (the promoter of maize polyubiquitin gene) will be used to drive the expression of the FLP gene, and the introduced DNA construct will contain the short version of the FRT sites.

 Although the site-specific recombination
10 sequences must be linked to the ends of the introduced DNA, the gene encoding the site specific recombinase may be located elsewhere. For example, the recombinase gene could already be present in the plant's DNA or could be supplied by a later introduced DNA fragment either introduced
15 directly into cells, or through cross-pollination.

 One method for targeting the insertion of a DNA sequence into the DNA of a eukaryotic cell involves the use of a specific DNA transformation construct. Various DNA sequences can be included as components of these DNA
20 transformation constructs, and are defined as follows. The "targeted DNA sequence" encompasses those nucleotide sequences present in the DNA construct, that are flanked by nucleotide sequences sharing homology to nucleotide sequences present in the eukaryotic cell (see Fig. 1A
25 (bracket a)). The "DNA sequence of interest" is a subset of the targeted DNA sequence, consisting of DNA sequences intended to be permanently inserted into the DNA of a eukaryotic cell. The "excisable selection region" is also a subset of the targeted DNA sequence, comprising DNA
30 sequences encoding a selectable marker flanked by a site-specific recombination sequence. The excisable selection region enables selection of homologously transformed cells, and the entire region can be removed after selection of cells transformed via homologous recombination. In
35 addition, the excisable selection region may also contain a

-12-

gene encoding a site-specific recombinase capable of interacting with the site-specific recombination sequence flanking the excisable selection region.

The selectable marker gene is operably linked to regulatory sequences capable of expressing the gene in the eukaryotic cell. The DNA sequence of interest is flanked by nucleotide sequences sharing homology with nucleotide sequences present in the eukaryotic cell. These flanking homologous sequences will induce a recombination event which inserts the targeted DNA region into a specific site of the eukaryotic cell's DNA.

The targeted DNA region of the transformation construct may comprise sequences encoding a selectable marker or a polylinker region. A polylinker is a short length of DNA that contains numerous different endonuclease restrictions sites located in close proximity. The presence of the polylinker is advantageous because it allows various expression cassettes to be easily inserted and removed, thus simplifying the process of making a construct containing a particular DNA fragment.

The above described transformation construct can also be part of a larger construct. The additional sequences of the larger construct comprising DNA sequences capable of replicating the entire DNA molecule in a bacterial host and DNA sequences encoding a bacterial selectable marker (such as genes encoding for ampicillin or tetracycline resistance). This larger construct, ideally a plasmid, can be used to transform bacterial cells. These transformed bacterial cells can then be cultured to produce large quantities of the plasmid DNA. The specific transformation construct can then be isolated using techniques well known to those familiar with the art.

Targeting the insertion of a DNA sequence into a specific site of a host cell's DNA requires introduction of the transformation construct into the cell by one of the

-13-

known transformation techniques. Once the DNA has entered the host cells, the cells can be cultured under a selection scheme that kills any cells not containing the selectable marker. After selection for transformed cells,

- 5 introduction of site specific recombinase activity will excise the randomly inserted DNA constructs. These excised DNA fragments will subsequently be lost from the cell via the action of nucleases. Those cells that have integrated the introduced DNA via homologous recombination will retain
10 the integrated DNA even in the presence of recombinase activity. These cells can subsequently be isolated and cultured to proliferate callus and regenerate new tissues, organs and organisms (plants or animals).

- 15 A recombinase gene cassette is used to generate the recombinase activity needed to excise the randomly integrated DNA fragments. This recombinase gene cassette can either be physically linked to the introduced DNA sequences, already present in the host cell's genome, or subsequently introduced as a separate DNA molecule.
20 However, the timing of recombinase expression relative to the integration process may be a critical factor in producing the desired end product. Homologous recombination takes place in both animal and plant cells, within approximately 30 minutes after transformation. An
25 additional 3-4 hours is required for the production of detectable amounts of foreign gene product. Applicants initially assume that upon co-introduction of the FLP gene and the transformation construct into protoplasts, integration and homologous recombination will be completed
30 prior to significant FLP recombinase activity. If the co-introduction strategy is too inefficient, strategies can be utilized that control the recombinase gene expression.

In one embodiment of the invention, recombinase activity is delayed for a sufficient time interval to allow

-14-

the gene of interest to be integrated into its specific site via homologous recombination.

One way of delaying the activity of the recombinase comprises operably linking the recombinase gene to an inducible promoter. Inducible promoters include any promoter capable of increasing the amount of gene product produced, by a given gene, in response to exposure to an inducer. Inducible promoters are known to those familiar with the art and a variety exist that could conceivably be used to drive expression of the recombinase gene.

Two preferred inducible promoters are the heat shock promoter and the glucocorticoid system. Promoters regulated by heat shock, such as the promoter normally associated with the gene encoding the 70-kDa heat shock protein, can increase expression several-fold after exposure to elevated temperatures. The heat shock promoter could be used as an environmentally inducible promoter for controlling transcription of the FLP gene. The glucocorticoid system also functions well in triggering the expression of genes including recombinase genes. The system consists of a gene encoding glucocorticoid receptor protein (GR) which in the presence of a steroid hormone forms a complex with the hormone. This complex then binds to a short nucleotide sequence (26 bp) named the glucocorticoid response element (GRE), and this binding activates the expression of linked genes.

The glucocorticoid system can be included in the DNA transformation construct as a means to induce recombinase expression. In one embodiment utilizing the glucocorticoid system, the entire introduced DNA is flanked by direct repeats of either the lox or FRT site-specific recombination sequence. Located between these direct repeats are several elements including: the DNA sequence of interest flanked by sequences sharing homology to host cell nucleotide sequences, the recombinase gene sequence (Cre or

-15-

FLP) driven by a truncated (and therefore ineffective) promoter containing at least one GRE sequence positioned upstream from the truncated promoter, a selectable marker gene driven by a constitutive promoter and the

5 glucocorticoid receptor (GR) gene driven by a constitutive promoter. Following transfer of this linear piece of DNA into the cell's nucleus, the gene of interest will be inserted into its specific site via homologous recombination. Multiple random insertions of the entire

10 piece of DNA will also occur throughout the genome. The resulting cells will then be cultured in the presence of a selection agent to isolate transformed cells. Once these cells/calli have been selected, they will then be treated with the steroid hormone, glucocorticoid or one of its

15 synthetic equivalents such as dexamethasone. The steroid hormone will associate with the constitutively produced GR protein to bind to the GRE elements, thus stimulating expression of the recombinase gene (eg. Cre or FLP). Production of the recombinase will then trigger the

20 excision of randomly inserted DNA sequences, but will not effect any insertion resulting from the homologous recombination event.

Inducible promoters are not the only means of triggering the production of recombinase activity. Other

25 potential means of delaying the functional activity of a gene product are known to those familiar with the art.

Once recombinase activity has been triggered and the randomly inserted DNA has been removed, various means can be used to isolate those cells/calli that contain the

30 DNA of interest. Using standard techniques known to those familiar with the field, eg. Southern blotting and polymerase chain reaction, DNA can be isolated from individual calli and screened for the presence of the DNA of interest. An alternative method of screening to

35 identify desired transformants utilizes a visible

-16-

phenotypic marker located on the homologously inserted DNA. Then only those calli or regenerated plants that express the phenotype have the desired DNA insertion.

In an alternative selection scheme the DNA sequences inserted via a homologous recombination event also include sequences encoding a selectable marker. This DNA transformation construct can be used to transfect cells, and transformed cells can be isolated by culturing the cells in the presence of selection agents. Once transformed cells have been isolated, and the randomly inserted DNA sequences have been removed, the cells can be exposed to a second round of selection. Only those cells transformed via a homologous recombination event will retain the selectable marker gene, and will survive the second round of selection and culturing.

In one embodiment, as shown in Figure 1A(A), a recombinant DNA molecule for use in transforming eukaryotic cells comprises a multifunctional DNA sequence flanked by two directly repeated site-specific recombination sequences. The multifunctional DNA sequence comprises a targeted DNA region flanked on each end by nucleotide sequences sharing homology to nucleotide sequences present in the eukaryotic cell to be transformed. The targeted DNA region comprises a selectable marker gene under the control of a constitutive promoter and a DNA sequence of interest (the DNA sequences desired to be inserted into the cells DNA).

Once the multifunctional DNA sequence is introduced into the cell, the entire molecule can be inserted into the host cell's DNA via a random recombination event, or alternatively the targeted DNA region can be inserted via a homologous recombination event. Selection for the expression of the selectable marker gene product will isolate both of these types of transformed cells. Subsequent site specific recombinase

-17-

activity, capable of interacting with the site-specific recombination sequences flanking the multifunctional DNA sequence, can then be generated within the cell. This recombina-
5 randomly inserted recombinant DNA molecules. Once the randomly inserted DNA sequences have been removed, exposure of the cells to a second round of selection for the selectable marker will isolate those cells containing the targeted DNA region integrated via homologous
10 recombination.

One manner of generating the site specific recombina-
se activity involves retransforming the cells with a second DNA construct that encodes a site specific recombina-
se. Alternatively, the recombina-
15 included on the original introduced DNA construct as shown in Figure 1A(B). If the site specific recombina-
se gene is co-introduced with the original DNA construct, the expression of the recombina-
se gene may be regulated by an inducible promoter. Potential inducible promoters include
20 the heat shock promoter and the glucocorticoid system.

As shown in Figure 1A(C), the transformation construct can also include a polylinker region located in the targeted DNA sequence. The addition of a polylinker region promotes the ease of constructing unique DNA
25 molecules. Through the use of specific nucleotide restriction enzymes, gene cassettes and other DNA sequences can be inserted into the polylinker region. Any DNA sequence inserted into the polylinker can then be targeted for insertion into a host cell's DNA via homologous
30 recombination.

In another embodiment, two different site specific recombination systems, eg., cre/lox and FLP/FRT are used in combination to select for cells transformed via homologous recombination. DNA sequences must first be
35 introduced into the host cells, using techniques known to

-18-

those familiar with the art. These cells will then be subjected to a first round of selection to isolate cells containing the introduced DNA integrated into the cell's DNA. This round of selection will yield a majority of
5 cells that have the introduced DNA randomly inserted into the cell's genome as well as a small number of cells having the introduced DNA integrated into the cell's genome by homologous recombination. It is highly probable that those
10 cells containing the introduced DNA integrated via a homologous recombination event will also have the introduced DNA randomly inserted at different locations of the genome. These randomly integrated sequences can be removed through the use of a site specific recombination system.

15 The second round of selection occurs after the randomly inserted DNA sequences have been removed from the host cell's genome. This second round of selection identifies those cells that have a portion of the original introduced DNA integrated into the predetermined site of
20 the host cell's DNA by homologous recombination. A second separate site specific recombination system can then be utilized to remove the DNA sequences that were required for the second round of selection, leaving only desired DNA sequences still integrated into the host cell's DNA.

25 To carry out this process the desired DNA sequences targeted for integration into the host cell's genome must be operably linked to a number of functional DNA sequences. First of all, any introduced DNA sequences that are likely to be randomly inserted into the host
30 genome must be flanked by two directly repeated site-specific recombination sequences. In this construction, any randomly integrated DNA sequence can be excised through the interaction of a site specific recombinase with its site-specific recombination sequence. A number of
35 different site specific recombinase systems can be used,

including the Cre/lox system of the bacteriophage P1, the FLP/FRT system of yeast, the Gin recombinase of phage Mu, and the R/RS system of pSR1 plasmid.

In one preferred embodiment the introduced DNA consists of a multifunctional DNA sequence flanked by direct repeats of a first site-specific recombination sequence (see Figure 1A). The DNA located between the two site-specific recombination sequences comprises, a gene encoding a first site specific recombinase gene capable of recognizing the first recombinase target sequence, and a targeted DNA sequence. The targeted DNA sequence (Fig. 1A (bracket a)) is flanked by nucleotide sequences sharing homology to nucleotide sequences present in the eukaryotic cells to be transformed. These "homologous regions" function to initiate and terminate a homologous recombination reaction that will insert the DNA sequences located between the two homologous regions into the host cell's DNA. Preferably, the DNA sequences located between the two homologous regions comprises a DNA sequence of interest, and a "excisable selection region. The DNA sequence of interest consists of DNA sequences that are desired to be permanently inserted into the host cell's DNA. The "excisable selection region" (Fig. 1A (bracket c)) contains DNA sequences encoding products that are required for selection of the homologous recombination event and is flanked by direct repeats of a site-specific recombination sequence.

The excisable selection region is no longer needed once the cells containing the homologous recombination event have been selected. Therefore, these sequences can be removed after the second round of selection through the use of a second site-specific recombinase system. Optimally, a different site-specific recombinase system is utilized than the one use to excise the randomly inserted DNA sequences. Thus in one form of

-20-

the invention, the excisable selection region is flanked by direct repeats of a second site-specific recombination sequence, and comprises a gene encoding a selectable marker, and a gene encoding a second site specific recombina-
5 recombina- gene capable of recognizing the second site-specific recombination sequence. An example of this embodiment is shown in Figure 1A(D). The entire second site-specific recombina- system is located within the targeted region. The DNA sequences that are desired to be
10 permanently inserted into the host cell's DNA are located outside the second recombina- system, but within the targeted region. In this transformation construct the expression of the second recombina- gene is preferably controlled by an inducible promoter.

15 Once the construct, shown in Figure 1A(D), is introduced into a cell, transformed cells can be selected through the use of the selectable marker gene. After the first round of selection, the expression of the first recombina- activity will result in the removal of
20 randomly inserted DNA sequences. Exposure of these cells to a second round of selection will isolate those cells which have the targeted region integrated into the cell's DNA via a homologous recombination event. These cells will contain the excisable selection region and the DNA
25 sequences of interest. Inducing the production of the second recombina- will result in the excision of the excisable selection region leaving only the DNA sequence of interest. Thus a transformed cell can be created that contains only the desired genes integrated into a
30 predetermined site of the host cell's DNA.

Although including the site-specific recombina- genes on the introduced DNA sequence is convenient, this is not necessary. These genes could be supplied on a separately introduced piece of DNA or could already be
35 present in the host cell's DNA or could be introduced by

-21-

cross pollination. Typically all the genes contained on the introduced DNA will be linked to regulatory sequences capable of expressing the gene's product in a eukaryotic cell. However, in preferred embodiments the expression of
5 at least one of the site specific recombinase gene will be regulated by an inducible promoter.

For those transformation constructs that allow direct selection of cells transformed via a homologous recombination event, only those cells containing DNA
10 integrated by homologous recombination should survive the second round of selection. However, the possibility does exist, because of some error in recombination, that the site specific recombinase activity will fail to remove all the randomly inserted DNA fragments. However, these
15 anomalies will normally be detected by standard genomic DNA analysis and can thus be eliminated.

As shown in Fig. 1B, constructs A - D can be physically linked to additional sequences to form a circularized DNA molecule (a plasmid). These additional
20 sequences would contain a gene cassette encoding a bacterial selectable marker and a bacterial origin of replication. This plasmid is useful in generating large amounts of the A - D DNA constructs. The procedure consists of using this plasmid to transform bacterial
25 cells, growing the bacterial cells under selection for the presence of the plasmid and then finally isolating the plasmid from the replicated bacterial cells. In order to prevent integration of unnecessary plasmid DNA that is outside the site-specific recombination sequences, the
30 plasmid will also contain restriction endonuclease sites that allow the removal of the plasmid sequences prior to transformation.

The following examples are intended to illustrate but not limit the invention. All plasmid constructs are
35 harbored in E. coli strain HB101 cultures maintained in our

-22-

laboratory under label corresponding to the plasmid construct designation.

EXAMPLE 1

5 Homologous Recombination between Plasmid DNAs

Materials and methods

Plasmid constructions. A 6.8 kb Bam HI-XhoI fragment of maize genomic DNA containing the *adh1*-S gene was inserted into the BamHI-SacI sites of pBR322 to yield pBx26. (See 10 Fig. 2A.) A control vector pAeiGUS, was constructed by replacing the PvuII fragment of pBx26 (containing most *adh1* coding sequences) with a SmaI-EcoRI fragment containing the *gusA* coding sequence and a *nos* polyA site. Thus, the construction contains a 1.5kb BamHI-PvuII fragment from the 15 *Adh1* genomic clone with the first exon and intron A of the *Adh1* gene fused in-frame to the *gusA* coding sequence; however, the *gusA* coding sequence still has its own start codon intact. The *Adh1* promoter was removed by elimination of the HindIII fragment from pAeiGUS to create piGUS. 20 piGUS still contained a 459 bp fragment of intron A in front of the *gusA* coding sequence, but the chimeric *gusA* gene was not active because of the absence of a promoter. The p35SGUS construction contained the GUS coding sequence driven by the CaMV 35S promoter. p35SiGUS was constructed 25 by insertion of the blunt-ended SacII-PvuII fragment of pBx26 containing intron A into the SmaI site of p35GUS. pAGUS was constructed by insertion of the BamHI-Csp45 promoter fragment of pBx26 into the BamHI-EcoRI site of pBI201.

30 Figs. 2A and B show plasmid constructions and general strategy of recombination experiments. (Fig. 2A) pBx26 (11.5 kb) was doubly digested with relevant restriction enzymes and *Eco*RI to standardize the 5' end of the DNA molecule participating in the recombination 35 reasction. This fragment (total size 2.1 kb) contained the

promoter (dotted bar), first exon (closed bar), and intron A (open bar) of the maize *Adh1* genomic clone. *piGUS* contained the *HindIII*-*PvuII* fragment of intron A followed by the β -glucuronidase (GUS) coding sequence and the
5 nopaline synthase (NOS) polyadenylation site. The *PvuII* site was destroyed after ligation to the GUS coding sequence. Only those restriction sites applicable to the studies reported here are shown. The *Adh1* coding sequence is not drawn to scale. (Fig. 2B) The effective homologous
10 region (*HindIII*-*PvuII* fragment of intron A) contained restriction sites for *BglI*, *BglII*, and *StuI* in the pBx26 and *piGUS* molecules. Numbers in parentheses represents the nucleotide position of the restriction sites on the pBx26 map.

15 All plasmid DNAs were prepared using the alkaline lysis method and purified by centrifugation in CsCl-ethidium bromide gradients as described in Maniatis et al. (1982). After the first centrifugation, the contaminating RNA was removed by incubation of the plasmid preparations
20 for 1 h at room temperature with DNase-free pancreatic RNase. Finally, plasmid DNA was phenol extracted and dissolved in TE(10mM TRIS-HCl, 1mM EDTA pH 8.0) buffer (at 1 mg/ml) according to standard procedures.

Cell culture, protoplast isolation, and transformation.

25 An A188xBMS cell suspension line of maize (*Zea mays* L.) was initiated from a type II friable callus as described by Kamo et al., *Planta* 172: 245-251 (1987). Suspension cultures were maintained on a gyratory shaker at 120 rpm and 25°C in the dark. The cells were subcultured
30 every week by placing 5ml packed cell volume into 35 ml of N6 medium (Chu et al. 1975) supplemented with 6 mM proline, 1 g/l casein hydrolysate, and 3.5 mg/l 2,4-dichlorophenoxyacetic acid (2,4-D). For protoplast isolation, 2 ml packed cell volume (pcv) were subcultured
35 for 3 to 10 days before isolation in 41 ml of MS medium

-24-

(Murashige and Skoog 1962) supplemented with 3.5 mg/l 2.4-D.

- Approximately 5 ml pcw suspension cells were digested for 3 h in 20 ml MS medium containing 0.2 M mannitol, 0.5 mg/l thiamine, 2 mg/l 2.4-D, 80 mM CaCl₂, 2% cellulase, 0.25% pectinase 0.1% (Worthington Biochemical Co.) pectolyase Y-23 (Seishin Pharmaceutical Co.) pH 6.0. The protoplasts were filtered through a 48 um screen and pelleted by centrifugation at 50 x g for 15 min. The protoplast pellet was suspended in 8 ml of protoplast culture medium (PCM); (Chourey and Zurawski, 1981, Theor Appl. Genet 59:341-344), containing 9% Ficoll (Ficoll 400, Sigma Chemical Co.), dispensed into two tubes, and overlaid with 4 ml of transformation solution (0.2M mannitol and 80 mM CaCl₂). After centrifugation at 75xg for 10 min, a band of protoplasts was collected at the interphase, and the concentration of protoplasts was adjusted to 1×10^7 protoplasts/ml with PCM. Transformation was carried out in 12 ml sterilized Falcon polystyrene tubes by pipetting an amount of plasmid DNA followed by 0.5 ml of protoplast suspension and 0.5 ml 50% PEG (polyethylene glycol, molecular weight 8000: Sigma Chemical Co.) dissolved in F-medium (Krens et al. 1982 Nature 296:72-74). Incubations were for 20 min at room temperature. After incubation, 0.33 ml of the protoplast suspension was transferred to each well of a 12-well microculture plate containing a 2 ml block of solidified 0.2% low melting point agarose (Bethesda Research Laboratories) in PCM; this procedure eliminated the need for manual dilution of PEG. Transformed protoplasts were incubated for 20-24 h at 26°C in the dark. DNA used for transformation was digested with appropriate restriction enzymes according to the recommendations of the manufacturers. The completion of the digestion and the concentration of DNA fragments were estimated by HPLC analysis of the digestion mixture on a

-25-

PE-TSK DEAEENPR column (Kato et al., 1989, Chromatography Journal, 478:204-268).

GUS activity determination. After incubation, protoplasts and all liquid from the cell culture were collected into 1.5 ml microcentrifuge tubes. After a brief centrifugation (5 s), the protoplast pellet was suspended in 0.3 ml GUS extraction buffer (Jefferson, 1987, Plant Mol Bio Rep 5:387-405) (containing 0.1% Triton X-100), mixed, and centrifuged. Protoplast extracts (0.1 ml) were incubated with 0.6 ml of 1 mM MUG (4 methyl umbelliferyl B-D-glucuronide) in GUS extraction buffer (without 0.1% Triton X-100) at 37° C. Three samples (0.2 ml incubation mixture in 1 ml of 0.2 M Ca₂CO₃) were used for fluorescence measurements (excitation at 365 nm, emission at 455 nm) in a Perkin Elmer spectrofluorimeter calibrated with standard solutions of methylumbelliferone. GUS activity was estimated from the slope of the line generated from time points and normalized to the protein content determined by the method of Lowry et al. (1951).

DNA analysis. Total DNA was isolated from 5x10⁶ protoplasts 20h after transformation. Protoplasts were incubated in PCM medium supplemented with 30 units of DNase for 1 h at 37°C to digest plasmid DNA that remained in solution. After incubation, protoplasts were centrifuged and the pellet was washed with PCM. DNA was isolated by repeated phenol-choloroform extraction of the protoplast pellet, and precipitated from the aqueous phase with 2 volumes of ethanol. After washing with 70% ethanol and vacuum drying, the DNA pellet was suspended in 10 ul of water. Five microliters of the DNA preparation were mixed with 50 nmol P1 primer (5'-GGGGTTTCTACAGGACG-3' (SEQ ID NO:1); specific for the GUS coding sequence) and 50 nmol P2 primer (5'-CCTCACAGGCTCATCTCG-3' (SEQ ID NO:2); specific for the Adhl promoter sequence), a dNTP mixture (final concentration 0.2 mM each), 5 ul of the 10x polymerase

-26-

chain reaction (PCR) buffer (Perkin-Elmer Cetus, Emeryville, Calif.), and Tag polymerase (2.5 units per tube). The MgCl₂ concentration of the buffer was adjusted to 2 mM, and the final volume of the reaction was adjusted to 50 ul with water. PCR was carried out in a Perkin-Elmer Cetus DNA thermal cycler. After initial heating at 94°C for 5 min, 25 cycles of PCR were carried out with denaturation, annealing, and extension steps of 94°C for 1 min, 45°C for 1 min, and 72°C for 2 min, respectively. The extension steps were increased with each cycle by an additional 15 s. Twenty-five microliters of the amplification reaction were loaded on an agarose gel to analyze the products.

For Southern blot analysis, DNA was isolated from transformed protoplasts as described above for PCR. The standard protocols of Maniatis et al., (1982) were used for the DNA electrophoresis, blotting, and hybridization. The radioactive probe was prepared using the Multiprime DNA labeling system of Amersham International according to the manufacturer's instructions. The blot was hybridized with a ³²P-labeled BamHI-SstI fragment of the pPUR plasmid (Lyznik, et al., 1989 - gusA coding sequence).

Results

Experimental system

The experimental strategy relied on the supposition that two separate DNA molecules with overlapping homologous regions, would recombine upon introduction into Maize protoplasts. The recombination event would result in the positioning of the gusA coding sequences behind the maize Adh1 promoter. The resultant GUS activity would indicate that homologous recombination had occurred between the two plasmids, since neither of the two separate plasmids could express gusA. Both plasmids, pBx26 and piGUS, contained 459 bp of Adh1 intron A to serve as the effective region of homology. Recombination within

this region of the plasmid would be expected to yield a functional gusA gene.

GUS expression in transformed protoplasts

The pAeiGUS plasmid was a much more effective
5 construction for GUS expression in maize protoplasts than
constructions based only on the Adhl promoter or the 35 S
CaMV promoter. Plasmid pAeiGUS (6.6kb) was about 1 kb
larger than p35SGUS (5.6 kb) and 0.4 kb larger than
p35SiGUS (6.2 kb); thus, it is probable that the higher GUS
10 expression from pAei-GUS was due to an increase expression
rate rather than increased uptake of DNA into the
protoplasts. GUS activity could be observed in maize
protoplasts 4 h after transformation with pAeiGUS. The 20-
24 h period between transformation and enzyme assay used in
15 experiments reported here was a compromise. This was owing
to the need to have a strong signal of GUS activity, but
also the need to keep the protoplast incubation time short.
Recombination reactions in mammalian systems take place
shortly after transformation, and the DNA persists in the
20 nucleus with a half-life of about 20 h. While variability
from experiment to experiment was quite high, variability
within a single experiment was low.

Evidence of homologous recombination

When maize protoplasts were co-transformed with
25 the products of the EcoRI-PvuII double digestion of pBx26
and HindIII-linearized piGUS DNA, GUS activity was observed
after the 20-24 h period of incubation. The transient
nature of GUS expression following the recombination
between free plasmid molecules was evident in the decline
30 in GUS activity over time. GUS activity was at the
background level after protoplast transformation with
plasmid DNAs supplied separately. These results indicated
that interaction of the two plasmid DNA molecules in the
co-transformed protoplasts was required for the detection
35 of GUS activity.

-28-

Support for the presence of recombinant DNA molecules in protoplasts was provided by PCR analysis of total DNA isolated from co-transformed protoplasts. The 720 bp DNA amplification product (see the Materials and methods for details of the two primers used) expected as a result of the recombination reaction was observed when the reaction was carried out in the presence of total DNA isolated from co-transformed protoplasts. The same sized fragment was amplified from total DNA isolated from protoplasts transformed with the control plasmid pAeiGUS, but not from DNA isolated from protoplasts following the treatments with piGUS, pBx26, or mock transformation (no DNA).

The amplification procedure of the polymerase chain reaction can produce recombinant molecules in vitro from piGUS and pBx26 DNA fragments sharing homologous sequences; thus, the observed 720 bp DNA fragment might only partially result from the amplification of recombination products generated inside the co-transformed protoplasts. However, direct analysis of total protoplast DNA on agarose gels followed by blotting and hybridization to a probe of the GUS coding sequence showed that DNA fragments of higher molecular weight than the original transforming DNA and containing the GUS coding sequences were present only in DNA preparations from co-transformed protoplasts. In addition, the linearized piGUS DNA molecules seemed to be more susceptible to degradation than circular pAeiGUS.

A simple ligation of free ends of DNA between restricted pBx26 and piGUS might lead to the activation of the *gusA* coding sequence. The DNA molecules did recombine (ligate) to activate the *gusA* gene, but at a very low level. The activity (close to the background level) represented only a small fraction of the GUS activity obtained after co-transformation with plasmid fragments

-29-

- sharing the overlapping effective homologous sequences. The ligation product of the co-transformed EcoRI-PvuII fragment of pBx26 and piGUS linearized with HindIII should yield a fragment longer (1.2 kb) than the 720 bp
- 5 amplification product obtained in the PCR. Indeed, the main amplification products following co-transformation of pBx26 and piGUS into maize protoplasts was of the size corresponding to the amplified fragment of intact pAeiGUS DNA with other only very minor bands seen in the original
- 10 get. GUS activity above the background level was observed after co-transformation with the circular forms of plasmid DNA or with plasmids linearized outside the effective homologous regions but was increased after restriction proximal to or inside the effective homologous regions.
- 15 These results indicated that the recombination between plasmid molecules was not a random process but, rather, depended strongly on both the lengths of the homologous regions and the positions of the cuts within the DNA sequences.
- 20 *Other requirements of recombination reactions*
- Both plasmid DNA molecules had to be linearized to obtain efficient expression of GUS protein; if only one of the plasmids was linearized the level of GUS activity, and hence recombination, was low. In addition, at least
- 25 one end of the linearized molecule had to share homology with the effective homologous region of the other molecule. The presence of a terminal nonhomologous region of the 3' end of the pBx26 EcoRI-PvuI or EcoRI-AccI fragments strongly reduced GUS activity. In all experiments
- 30 performed involving pBx26 and piGUS, the highest GUS activity was observed in protoplasts co-transformed with the EcoRI-PvuII fragment of pBx26. This fragment contained the maximum length (459 pb) of the effective homologous region.

-30-

Similar digestions of the other recombination substrate, piGUS, did not provide conclusive results. There was no difference in the total length of the effective homologous region between products of different restriction enzyme digestions since all restriction enzymes used piGUS cut only in one place. While HindIII digestion pf piGUS produced the highest GUS activity when co-transformed with the pBx26 EcoRI-PvuII fragment, in other experiments StuI-digested piGUS molecules were more efficient. StuI digestion resulted in 113 bp of the effective homologous region at the 5' end of the piGUS DNA as compared to the 459 bp after HindIII digestion. Since the enzyme only cut the plasmid at one site, the products of the StuI restriction contained homologous regions at both ends of the DNA molecule; HindIII digestion resulted in the effective homologous region being only at one end.

Double digestion of piGUS DNA with StuI-BglII and StuI-HindIII or StuI-PvuII; (removal of increasing lengths of the 3'-end terminal homology) resulted in decreased activity of GUS in co-transformed protoplasts as compared to the single digestion with StuI. This effect was observed when the digestion products were separated before co-transformation or when a mixture of restriction fragments was used in co-transformation experiments. Thus, it was unlikely that the observed decrease in GUS activity was due to the interference of the additional digestion products (StuI-BglII or StuI-HindIII short fragments) with the recombination reaction. Once the 3' end of piGUS did not contain a homologous region (HindIII digestion), additional removal of plasmid sequences at this end did not affect the recombination activities.

Discussion

It is shown that homologous recombination occurs between two plasmids following their uptake into maize protoplasts. Use of the *gusA* reporter gene, the transient

expression assay for GUS, and the direct uptake of DNA mediated by PEG made the experimental system rapid, convenient, and reliable. The use of two plasmids, each incapable of expressing *gusA* alone, but sharing homologous sequences provided by intron A of the *adh1* gene of maize, permitted the direct demonstration of homologous recombination through the expression of *gusA*. By restricting the homologous regions of the two plasmids at different sites, information was obtained concerning recombination reactions. This study has shown that homologous recombination between introduced plasmid DNA can take place in plant protoplasts.

EXAMPLE 2

FLP Mediated Site Specific Recombination between Plasmid FRT sites in Maize and Rice cells

Materials and Methods

Synthesis of FLP expression vectors

The genomic clone of maize *adh 1* and the vector of pAeiGUS were described in Example 1. Plasmid pAHC27 was provided by Dr. Peter Quail, University of California, Berkeley. Plasmids pNEO β GAL and pOG44 were purchased from Stratagene, LaJolla, CA. pUbiGUS was constructed by ligating the maize ubiquitin gene 5' controlling elements in front of the *gusA* coding sequence by ligation of the XbaI fragment of pAHC27 into HindIII-BamHI restriction sites of p35SGUS. For the construction of pAeiFLP, both pAeiGUS and pOG44 were digested with *Bgl*III and *Sac*I. The 1.5 kb fragment carrying the FLP coding sequence from pOG44 was then ligated directly into the eluted 4.4 kb fragment of pAeiGUS replacing the *gusA* coding sequence. pUbiFLP was obtained by isolating the 1.5 kb *Sac*I-HindIII FLP fragment from pOG44 and subcloning into the respective sites of pGEM-7Z(-)f (Promega, Madison, WI)). The resulting plasmid was cleaved with *Sac*I and SmaI, and the 1.5 kb FLP fragment

-32-

was then ligated directly into *SacI*-*SmaI* digested pUbiGUS replacing the *gusA* fragment with the FLP gene.

Synthesis of FRT containing vectors

Two primers

- 5 (5'-GTGATCAGAAGTTCCTATTCCGAAGTTCCTATTCTCTAGAAA-3' (SEQ ID NO:3)) and (5'-CTGATCAGAAGTTCCTATACTTTCTAGA-3' (SEQ ID NO:4)) were annealed (4nmoles each) and incubated with 5 units of T4 DNA polymerase and 60 nmoles of each dNTP in 0.1 ml at 11°C for 3 hrs. to form a complete FRT
- 10 recombination site of 48 bp. The primer-extended fragments contained a *BclI* restriction site on each end. Phosphate groups were then added to the terminal nucleotides by incubation with 20 units of T4 DNA kinase and 16 nmoles ATP in 30 μ l of 1x kinase buffer. The resulting
- 15 double-stranded DNA fragments were blunt-end ligated and cleaved with *BclI* restriction enzyme. The products were then ligated directly into the *BglIII* site of pUbiGUS forming pUFRTG. For the construction of the pUFRTmG vector, containing a minimal version of the FRT site (37
- 20 bp), the primer extended products were blunt-end ligated into the *BglIII* site of pUbiGUS (the *BglIII* site of pUbiGUS was filled-in using T4 DNA polymerase). Confirmation of the minimal FRT sequence was done by sequencing the double-stranded plasmid DNA. pU2FRTG and pU2FRTmG were
- 25 constructed by ligation of the 1.3 kb *XbaI* fragment of pNEOBGAL into pUFRTG and pUFRTmG, respectively. The test plasmid pUbiFRT for intermolecular recombination was constructed by *EcoRI* restriction of pU2FRTmG and religation of the resulting plasmid DNA fragments. The other
- 30 substrate for intermolecular recombination, pFRTGUS, was obtained by digestion of pU2FRTmG with *EcoRI* and *BamHI*, isolation of the fragment containing the *gusA* coding sequence and the FRT site, and ligation of the resulting fragment into the respective sites of pGEM-7Z(-)f (Promega,
- 35 Madison, WI).

-33-

Molecular analysis of DNA

Polymerase chain reactions (PCRs) were carried out in a Perkin-Elmer Cetus DNA thermal cycler.

Amplification was performed using reagents and protocols as

- 5 outlined by the Perkin-Elmer Cetus GeneAmp PCR Kit (Perkin-
elmer Cetus, Norwalk, CT). Sequences of primers were as
follows: 5'-CCCCAACCTCGTG-3' (SEQ ID NO:5) for the first
exon of the ubiquitin gene and 5'-GGGGTTTCTACAGGACG-3' (SEQ
ID NO:1) for the 5' end of the *gusA* coding sequence. PCR
10 reactions contained: 5 μ l of template DNA, 5 μ l of primer
solution (50 mmoles each), 5 μ l of 10x PCR buffer II (22),
2.0 μ l dNTPs mix (final concentration 200 μ M of each
nucleotide), and 0.25 μ l of Taq polymerase (2.5
units/100 μ l) in a final volume of 50 μ l. Denaturation,
15 annealing, and extension steps were performed at 94°C for
1 minute, 45°C for 1 minute, and 72°C for 2 minutes,
respectively. The extension steps were increased 15
seconds with each cycle. Five microliters of each
amplified product were analyzed using 1.0% agarose gel
20 electrophoresis.

Southern blotting of agarose gels containing PCR
amplified DNA was performed using capillary transfer to
Hybond-N membrane (Amersham, Arlington Heights, IL). DNA
was fixed to the membrane by UV irradiation and incubated
25 in prehybridization solution (5x SSPE, 5x Denhardt's
solution, 0.5% SDS) at 65°C for 4 hours. The radioactive
probe (α ³²P-labeled *Sma*I- *Bgl*III fragment isolated from
pUbIGUS) was prepared using the Multiprime DNA labeling
system according to the manufacturer's instructions

- 30 (Amersham, Arlington Heights, IL). The probe contained the
complete ubiquitin intron and part of the first exon. The
blot was hybridized overnight in prehybridization solution
at 65°C and subsequently processed according to standard
procedures.

-34-

The nucleotide sequences of the FRT sites were determined by a modified dideoxy method using Sequenase Version 2.0 (United States Biochemical Cleveland, OH). CsCl purified pUFRTG and pU2FRTmG plasmid DNA was used as template DNA. The primer used was the same as that used for PCR which annealed to the 5' end of the first exon of the ubiquitin gene.

Transformation procedures

The cell suspension culture of maize (*Zea mays* L.) was initiated from A188 x BMS type II callus and maintained (Kamo et al. 1987, Planta, 172, 245-251). Seven days prior to protoplast isolation, 2 ml packed cell volume (PCV) of the suspension culture was transferred into 37 ml of MS medium (Murashige et al., 1962, Physical Plant, 15, 473-497) supplemented with 3.5 mg/l 2,4-D.

Approximately 5 ml PCV of suspension cells were digested for 3 h in 20 ml MS medium containing 0.2M mannitol, 0.5 mg/l thiamine, 2 mg/12,4-D, 80 mM $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$, 2% cellulase, 0.25% pectinase, and 0.1% pectolyase Y-23, pH 6.0. Protoplasts were filtered through a 48 μm nylon mesh screen and pelleted by centrifugation at 50 x g for 15 minutes. The pellet was suspended in 8 ml of protoplast culture medium (PCM containing 9% Ficoll 400 and overlaid with 4 ml of transformation medium (TM)). The transformation medium consisted of 100 mM MES buffer, pH 5.5, 0.2 M mannitol, and 80 mM $\text{CaCl}_2 \cdot \text{H}_2\text{O}$. Following centrifugation at 75 x g for 10 minutes, a band of protoplasts was collected from the interphase, and the concentration was adjusted to 1.0×10^7 protoplasts/ml with TM.

Protoplast transformation was performed in 12 ml Falcon polystyrene tubes by pipetting 20-25 μl of plasmid DNA (1.0 mg/ml) followed by 0.5 ml of protoplasts and 0.5 ml of 50% PEG (polyethylene glycol, MW=8000; dissolved in F-solution). Protoplasts were incubated for 20 minutes at

-35-

room temperature. After incubation, 330 μ l of the protoplast solution were transferred to a 12-well microculture dish containing 2 ml solidified 0.8% low melting point agarose in PCM. Plates were then wrapped with Parafilm and incubated at 25°C in the dark for approximately 24 hours.

GUS Activity and Protein Determination

Following the 24 hrs. incubation, protoplasts were collected and resuspended in 300 μ l of GUS extraction buffer containing 0.1% Triton X-100. After centrifugation at 16,000 x g for 5 minutes, a 25 μ l extract of the protoplasts was incubated with 150 μ l of 1mM MUG (4-methylumbelliferyl β -D-glucuronide) in GUS extraction buffer in a 96-well plate at 37°C. Reactions were stopped at various times by adding 125 μ l of GUS stop buffer (0.2 M CaCO_3). Fluorescence (excitation at 362 λ and emission at 455 λ) was measured in a Perkin Elmer Luminescence Spectrometer LS50B calibrated with standards of methyumbelliferone. GUS activity was calculated from the slope of the line generated from time points and normalized to the protein content determined by the method of Bradford, 1976, Anal. Biochem., 72, 248-254.

Results

The FLP/FRT site-specific recombination system used in these studies consists of two elements: plasmid DNA encoding for the FLP enzyme and test plasmids containing the FRT recombination sites (Figs. 3A-B and 4A-B).

The target FRT nucleotide sequence consists of three recreated DNA sequences of 13 bp each; two repeats in a direct orientation and one repeat inverted relative to the other two. In addition, there is an 8 bp spacer region between the repeats which determines the orientation of the FRT recombination site and which serves as the site of sequence pairing that initiates the recombination event.

-36-

The FLP recombinase binds to all three repeats, anneals DNA sequences within spacer regions of two FRT sites, cleaves the sites at the borders of the spacer, and exchanges the DNA strands. Depending on the orientation of the recombination sites, the DNA fragment between the FRTs can be either inverted or excised. Additionally, the FLP recombinase can act on target FRT sites located on separate DNA molecules. These intermolecular recombinations can lead to integration of foreign DNA into FRT sites in bacterial and mouse genomes.

Fig. 3 illustrates the strategy and diagram of the FLP expression vectors used to study activity of yeast FLP/FRT system in plant protoplasts. More particularly Fig 3A shows components of the recombination system to test the DNA excision reaction catalyzed by the FLP protein. FLP enzyme can be produced by either pAeiFLP or pUbiFLP. Fig. 3B shows the sequence of the splicing site of the original *Adh1* genomic clone (SEQ ID NO:6) and the sequence of the junction sites between cloned FLP coding sequence and *Adh1* (SEQ ID NO:7) or ubiquitin maize (SEQ ID NO:8) promoters. Thick lines indicate an open reading frame of *Adh1* gene or the translation start codon for FLP protein synthesis in the pUbiFLP vector. Note that the first exon of the maize ubiquitin gene in contrast to the *Adh1* first exon is not translated.

Fig. 4 illustrates the structure of the FRT site-containing vectors. A single complete (SEQ ID NO:9) or partial (SEQ ID NO:10) FRT site was ligated into the *Bgl*III site of the ubiquitin first exon. Asterisks show the FLP binding sites. Arrows denote 13 bp inverted repeats. pUFRTmG vector contained only one 13 bp repeat and one shorter 11 bp inverted repeat. This FRT site lacks additional five FLP protein binding sites. Insertion of the *Xba*I fragment of pNEO β GAL into the *Xba*I site of the FRT

-37-

vectors provided the second FRT site and led to inactivation of GUS expression.

The FLP recombinase expressed from FLP vectors should recombine test plasmid DNA within the FRT target sites to restore expression of GUS enzyme. Expression of the FLP gene was driven by the maize *adh1* promoter in pAeiFLP and by the maize ubiquitin promoter in pUbiFLP (Fig. 3B). Both constructs included the first exon and intron of the respective genes. In the pAeiFLP construction, the first intron contained 290 bp of the first *adh1* intron fused at the *Bgl*III site to 81 bp of the synthetic intron from pOG44. Both promoters proved to be very effective as determined in transient GUS expression assays of maize protoplasts; however, the ubiquitin promoter was superior.

The *Bgl*III site of the first exon of the ubiquitin promoter in pUbiGUS was chosen for insertion of the FRT sequences following initial screening of other appropriate insertion sites. Two different FRT sites were ligated into the *Bgl*III site - one minimal 37 bp FRT site referred to as FRT_m and one complete 48 bp FRT site referred to as FRT (Fig. 4A). The FRT_m site lacks the third repeat which includes five additional FLP protein binding sites; however, a similar FRT deletion (missing one repeat and the same FLP protein binding sites) has been shown to be as effective as the wild-type FRT site in vitro. Interestingly, insertion of the FRT_m site into the pUbiGUS resulted in higher expression of the GUS enzyme. Insertion of the full length FRT site resulted in reduction of GUS expression.

A second FRT site for intramolecular recombination of test plasmids was created by insertion of the *Xba*I fragment of pNEO β GAL into the single *Xba*I site of pUFRTG. Insertion of the pNEO β GAL fragment provided supplemental sequences to form an additional FRT site and a

-38-

1.31 kb spacer DNA which separated the ubiquitin promoter from the *gusA* coding sequence (Fig. 4B). As a result, GUS activity in maize protoplasts transformed with pU2FRTG was substantially reduced. The reverse orientation of the

5 ligated *Xba*I fragment of pNEO β GAL should also inactivate GUS expression but should not form functional FRT sites. pU2FRTmG rev. and pU2FRTG rev. vectors were used to demonstrate that fully functional FRT sites were required to obtain activation of GUS expression by the FLP protein.

10 Transient GUS activity in maize protoplasts co-transformed with pAeiFLP and pU2FRTmG was higher than in protoplasts transformed with only pU2FRTmG. The restored GUS expression was approximately 10% of that observed after protoplast transformation with the control pUFRTmG. GUS

15 activity in maize protoplasts transformed with pAeiFLP was similar to background GUS activity. Thus, the FLP/FRT recombination system appeared to function in maize protoplasts. This was further substantiated by the increase of GUS activity in maize protoplasts co-

20 transformed with increasing amounts of pAeiFLP DNA. This indicated not only that GUS expression is dependent on the amount of the FLP enzyme present, but also that the amount of the FLP protein might have been the limiting factor in the recombination process. When pUbiFLP was introduced

25 into maize protoplasts along with the test plasmids, GUS activities were restored for both pU2FRTmG and pU2FRTG to 81% and 45% of the respective control treatments. Reactivation of GUS in the presence of pUbiFLP and the vectors containing both FRT's indicated that FLP protein

30 catalyzed excision of the 1.31 kb fragment. When the FRT's were mutated by the reverse ligation of the 1.31 kb fragment (pU2FRTmG rev. and pU2FRTG rev.), GUS activity was not restored indicating that these FRT sites were not recognized by the FLP protein.

-39-

To observe intermolecular recombination at the FRT sites, the ubiquitin promoter or the *gusA* coding sequence was removed from pUFRTmG to form pUbiFRT and pFRTGUS, respectively. Introduction of two different FRT sites into protoplasts on two separate DNA molecules (pUbiFRT and pFRTGUS) produced approximately a 4-fold increase in GUS activity above background activity in protoplasts transformed with pFRTGUS alone. This activity was 15% of the positive control (protoplasts transformed with pUFRTmG). The relatively high background expression of GUS in pFRTGUS transformed protoplasts (6 ± 1 unit) might be the result of the entire first intron and part of the ubiquitin first exon being present in this vector.

The FLP/FRT recombination system has also been tested in rice protoplasts. The recombinase-mediated excision of DNA proved to be as effective in rice as in maize protoplasts. The same pattern of response was observed for the different plasmid constructions containing FRTs. Co-transformation of rice protoplasts with pUbiFLP and either pU2FRTmG or pU2FRTG gave 75% and 31% restoration of the control GUS activity. Vectors containing inactive FRT sites gave only 1-2% of the control GUS activity.

PCR analysis of total DNA isolated from co-transformed maize protoplasts indicated the presence of recombinant plasmid DNA molecules. The two primers used for this analysis amplify the region between the ubiquitin transcription start and the 5' end of the *gusA* coding sequence. The length of this region is 2.49 kb in pU2FRTmG and pU2FRTG. After removal of the 1.31 kb fragment as a result of the recombination reaction, the amplified region should be reduced to 1.18 kb and should then be equivalent to the amplification product of pUFRTG DNA. A 1.18 kb fragment was identified in the products of the PCR reaction using total DNA isolated from maize protoplasts co-transformed with pU2FRTmG and pUbiFLP as a template. The

-40-

identity of this fragment was confirmed through hybridization using a probe specific to the first intron of the ubiquitin gene.

Discussion

5 The results presented here show that the FLP recombinase of yeast can promote the site-specific recombination between FRT sites in both maize and rice cells. This conclusion is based on the following evidence. Transient expression of the GUS enzyme from recombination
10 test vectors increased in protoplasts co-transformed with functional FLP expression vectors, and the magnitude of this increase depended on the amount of co-transformed plasmid DNA containing the FLP gene. Additionally, higher expression of GUS protein in co-transformed protoplasts was
15 observed when a stronger promoter was used to drive expression of the FLP protein. The increase in GUS expression was abolished if the FRT sites in test vectors were mutated. Lastly, the products of the site-specific recombination reaction were identified in co-transformed
20 protoplasts.

EXAMPLE 3

Integrative Recombination

To test the efficiency of insertion recombination
25 reactions, DNA constructs containing an FRT site integrated into a selectable marker gene are designed and synthesized. Upon introduction of these constructs into maize protoplasts followed by the selection of transformed calli, only material containing the integrated FRT sites is
30 selected. The neomycin phosphotransferase neo gene is used as a selectable marker and the 5' controlling elements (promoter) of the maize polyubiquitin gene is used to make pUFRTNeo (Fig. 5).

The neo gene has proven to be a reliable
35 selectable marker for maize protoplasts, and the ubiquitin

-41-

promoter assures a high level of foreign gene expression in transformed maize protoplasts. The coding sequence of the neo gene is not modified, but instead, the FRT site is ligated into the first untranslated exon of the ubiquitin leader sequence. This modification of the first exon does not abolish activity of the ubiquitin promoter.

Neomycin phosphotransferase activity and the copy number of integrated neo genes are determined in selected calli. Individual callus are selected (each containing a different number of active neo gene(s)) and used to establish suspension cultures.

Protoplasts isolated from these suspension cultures are re-transformed with DNA constructs containing a second FRT site and an inactive selectable marker. A plasmid DNA construction containing a promoterless bar gene that retains the first exon and intron of the maize ubiquitin leader sequence and an integrated FRT sequence (pFRTbar) is utilized. Re-transformed protoplasts are selected on medium containing phosphinothricin (PPT); only cells expressing the bar gene inactivate PPT and survive. The FLP recombinase activity is provided by co-transformation of protoplasts with the pUbiFLP plasmid DNA (described in Example 2). As a negative control, protoplasts are re-transformed with a construct similar to the pFRTbar construct except lacking the FRT site. Results are adjusted for activation of the bar gene due to spurious random integrations or homologous recombination within the ubiquitin first intron sequences.

Phosphinothricin-resistant calli are obtained as a result of FLP catalyzed site-specific recombination between FRT sites which places the ubiquitin promoter in close proximity to the bar gene. The efficiency of this reaction is analyzed by at least two methods. For calli containing a single or a few genomic DNA integrated FRT sites, the number of PPT-resistant calli produced after re-

-42-

transformation with pFRTbar versus the number of PPT-resistant calli obtained after retransformation with the pFRTbar construct lacking a FRT site, should indicate the efficiency of the site specific recombination process.

- 5 Results are also compared to the transformation efficiency of protoplasts transformed with a DNA construct containing the active *bar* gene (pUFRTbar), and to the co-transformation efficiency in the presence and absence of the pUbiFLP construct (to test the requirement of the FLP recombina
- 10 se activity).

- For protoplasts containing many copies of the FRT sites, an alternative method for assessing the recognition efficiency of the FLP recombina
- 15 se for genome-integrated FRT sites involves the analysis of *bar* gene integration patterns. This method requires the selection of re-transformed protoplasts on PPT medium, which results in both *bar*⁺ *neo*⁻ phenotypes (if all *neo* genes are inactivated) and *bar*⁺*neo*⁺ phenotypes (if only some of the *neo* genes are inactivated). All PPT-resistant calli can be
- 20 conveniently screened for *neo* activity using a recently modified microplate assay (Peng. et al., 1993, *Plant Mol Biol Rep*, in press). It is not necessary to make adjustments for re-transformation or co-transformation efficiencies since the product of a single transformation
- 25 event is being analyzed. The fraction of integrated FRT sites recognized as a site of *bar* gene integration indicates the effectiveness of the recombination reaction per se as well as the effectiveness of the communication network between DNA sequences inside the plant nucleus.
- 30 Several primary FRT transformed callus cell lines are tested to assess the influence of genomic position of the FRT sites on the recognition efficiency of integrated FRT sites by the FLP recombina
- 35 se. Site-specific integrations of the *bar* genes are verified and analyzed by Southern blotting and PCR.

EXAMPLE 4

Excisional Recombination

- One goal of the excisional recombination experiments is to determine the maximum distance between the two FRT sites that will still allow efficient excision by the recombinase. This maximum distance is determined through the use of the pU2FRTGUS DNA construct. This DNA construct contains a 1.3 kb DNA fragment with a unique BamHI restriction site located between the two FRT sites.
- 10 The BamHI restriction site allows for the insertion of additional DNA sequences to increase the distance between the two FRT sites. The efficiency of the excision reaction, relative to the distance between the two FRT sites, is evaluated using the transient GUS expression
- 15 assay.

Two different methods can be used to evaluate the efficiency of the FLP recombinase in the excision of integrated foreign genes: retransformation and controlled expression of the recombinase gene.

20 **Re-transformation**

- In the first method, maize protoplasts are transformed with a DNA construct containing the gusA gene inactivated by insertion of the neo gene into the promoter region of the gusA gene (pUFRTNeoFRTGus). The neo gene is
- 25 positioned between two directly-repeated FRT sites and is expressed (Fig. 6).

- Selection of transformed protoplasts on medium with kanamycin results in transgenic calli with a neo+ gus- phenotype. These calli serve as initial material to
- 30 establish new suspension cultures. Protoplasts isolated from these suspension cultures and re-transformed with the pUbiFLP DNA construct excise the neo gene and thus yield gus+ phenotypes, which can be detected in transient assays. These relatively simple experiments provide information

-44-

about the existence of the excision reaction catalyzed by the FLP protein in plant cells.

Controlled Expression of the FLP Gene

In the second approach, expression of the FLP gene is controlled to achieve excision so as to eliminate the re-transformation step in favor of controlling the FLP recombinase activity. Maize protoplasts are co-transformed with the pUFRTNeoFRTGus plasmid DNA and plasmid constructions containing the FLP gene under control of the soybean heat shock inducible promoter. Studies of the activity of the soybean Gmhsp 17.5-E gene promoter in maize protoplasts and cells showed excellent performance of this promoter in response to heat shock treatment. The background activity at 24°C was low and a several fold increase of activity was observed after heat shock at 42°C for 0.5-1 hour. The DNA construct containing the FLP gene under the control of this heat shock promoter was made (pHSFLP) and tested in transient assays for site-specific recombination (Table 1).

TABLE 1

Vector used for transformation	GUS activity nmol MU/min/mg protein			GUS activity corrected for pU2FRTmG background			GUS activity corrected for pUFRTmG expression			GUS activity corrected for pUbiFLP expression		
	Heat shock (hrs)			Heat shock (hrs)			Heat shock (hrs)			Heat shock (hrs)		
	0	0.5	1	0	0.5	1	0	0.5	1	0	0.5	1
-DNA	0.06											
pUFRTmG	58	41	41	58	41	41	58	58	58	58	58	58
pU2FRTmG	0.6	0.6	0.4	0	0	0	0	0	0	0	0	0
+ pUbiFLP	58	33	20	58	33	20	58	47	28	58	58	58
+ pHSFLP	0.8	8.4	9.1	0.2	7.8	8.7	0.2	11.1	12.4	0.2	14.0	24.8

There still remains the possibility that the residual FLP activity at room temperature might be too high, which could lead to excision reactions prior to the heat shock treatment. Modifications to the transformation procedure are made as necessary, such as growing the transformed protoplasts and cells at lower temperatures.

-45-

Once stably transformed calli are selected on medium containing kanamycin, the heat shock treatment activates the FLP gene leading to the excision of the neo gene and activation of the gusA gene. Three different phenotypes are generated: neo+gus-, neo+gus+, neo-gus+. The relative frequency of these phenotypes in relation to the copy number of integrated plasmid molecules provides an estimate of the efficiency of the excision reaction.

10

EXAMPLE 5

Gene Targeting and Excision of Foreign DNA

A strategy is developed as illustrated in Fig. 7 to use the FLP/FRT system to positively select from gene targeting events and at the same time eliminate any randomly integrated foreign plasmid DNA.

15

The adh1 locus of the maize genome is well characterized. The adh1 locus is used as a target site. DNA fragments of the adh1 genomic clone, about 5kb in length, is ligated between two directly-repeated FRT sites.

20

Gene targeting experiments on animal systems indicate that the efficiency of gene targeting increases exponentially with increasing length of the homologous regions on plasmid DNA molecules, but reaches a saturation at about 4 kb. A plasmid construct is made in which an active neo gene with

25

its own promoter (the CaMV 35S promoter) is inserted into the cloned adh1 homologous region. The second element of the site-specific recombination system - the FLP gene - is positioned outside the plasmid DNA region encompassed by the FRT sites. A schematic drawing of the targeting DNA

30

construct is presented in Fig. 8.

35

Maize protoplasts isolated from a A188xBMS suspension culture are transformed with the targeting DNA construct using the PEG-mediated transformation technique. The plasmid DNA is linearized outside the adh1 homologous region to produce a sequence replacement rather than a

-46-

sequence insertion DNA construct. Transformed protoplasts are selected on medium with kanamycin.

- DNA of the sequence insertion construct integrates into the genomic DNA via homologous recombination, and kanamycin resistant calli are recovered. These double reciprocal recombination events lead to replacement of the genomic adh1 DNA sequences with the neo gene of the plasmid DNA. Alternatively or simultaneously, the entire plasmid molecule integrates randomly through a random recombination mechanism. Any random integration event leads to expression of the FLP gene thus causing excision of the randomly integrated neo gene. Excision of the neo gene renders those cells sensitive to kanamycin and they subsequently die. Only those cells containing the neo gene residing within the genomic adh1 locus, as a result of homologous recombination, retain the resistance to kanamycin, and survive. The selected kanamycin-resistant calli are recovered and the DNA insertion events are analyzed by Southern blotting and/or the polymerase chain reaction to verify and demonstrate the occurrence of the targeting events.

- The FLP gene is positioned outside the DNA sequences flanked by the FRTs. This is due to the concern that the efficiency of excisions may be compromised if the FRT sites are separated by too great a distance. The FRT sites are separated by approximately 5.0 kb of DNA sequence in yeast 2 μ m plasmid DNA. This natural configuration of the FRT sites for the FLP activity should be considered as an acceptable first approximation. The transient assays for testing the efficiency of excision yield information about the flexibility of the FRT/FLP recombination system with regard to the distance between the FRT sites in maize cells. Where larger distances are tolerated by the recombinase, then the FLP gene and other components can be included with the DNA sequences flanked by the FRT sites.

-47-

Fig. 9 illustrates several examples of vectors in accordance with this invention adapted to control gene targeting and gene excisions using recombinases and inducible promoters.

-48-

SEQUENCE LISTING

(1) GENERAL INFORMATION:

- (i) APPLICANT: Hodges, Thomas K.
5 Lyznik, Leszek A.
- (ii) TITLE OF INVENTION: Controlled Modification of
Eukaryotic
Genomes
- (iii) NUMBER OF SEQUENCES: 10
- 10 (iv) CORRESPONDENCE ADDRESS:
(A) ADDRESSEE: Barnes & Thornburg
(B) STREET: 11 South Meridian St.
(C) CITY: Indianapolis
(D) STATE: Indiana
15 (E) COUNTRY: U.S.A.
(F) ZIP: 46204
- (v) COMPUTER READABLE FORM:
(A) MEDIUM TYPE: Floppy disk
(B) COMPUTER: IBM PC compatible
20 (C) OPERATING SYSTEM: PC-DOS/MS-DOS
(D) SOFTWARE: PatentIn Release #1.0, Version
#1.25
- (vi) CURRENT APPLICATION DATA:
(A) APPLICATION NUMBER: US 08/010,997
25 (B) FILING DATE: 29-JAN-1993
(C) CLASSIFICATION: 435
- (viii) ATTORNEY/AGENT INFORMATION:
(A) NAME: Lammert, Steven R.
(B) REGISTRATION NUMBER: 27,653
30 (C) REFERENCE/DOCKET NUMBER: 08/010,997
- (ix) TELECOMMUNICATION INFORMATION:
(A) TELEPHONE: (317) 231-7258
(B) TELEFAX: (317) 231-7433
- (2) INFORMATION FOR SEQ ID NO:1:
35 (i) SEQUENCE CHARACTERISTICS:

-49-

- (A) LENGTH: 17 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- 5 (ii) MOLECULE TYPE: DNA (genomic)
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:
- GGGGTTTCTA CAGGACG
- 17
- (2) INFORMATION FOR SEQ ID NO:2:
- 10 (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 18 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- 15 (ii) MOLECULE TYPE: DNA (genomic)
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:
- CCTCACAGGC TCATCTCG
- 18
- (2) INFORMATION FOR SEQ ID NO:3:
- 20 (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 42 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- 25 (ii) MOLECULE TYPE: DNA (genomic)
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:
- GTGATCAGAA GTTCCTATTC CGAAGTTCCT ATTCTCTAGA AA
- 42
- (2) INFORMATION FOR SEQ ID NO:4:
- 30 (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 28 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- 35 (ii) MOLECULE TYPE: DNA (genomic)

-50-

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:
CTGATCAGAA GTTCCTATAC TTTCTAGA

28

(2) INFORMATION FOR SEQ ID NO:5:

- 5 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 13 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

10 (ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:
CCCCAACCTC GTG

13

(2) INFORMATION FOR SEQ ID NO:6:

- 15 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 20 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

20 (ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:
CCCGTGCAGC TGC GGTGGCA

20

(2) INFORMATION FOR SEQ ID NO:7:

- 25 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 26 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

30 (ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:
CAACTGCAGC CCAAGCTTCC ACCATG

26

(2) INFORMATION FOR SEQ ID NO:8:

- 35 (i) SEQUENCE CHARACTERISTICS:

-51-

- (A) LENGTH: 56 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

5 (ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

CTTCTGCAGG TCGACTCTAG GATCCCCGGG TTCGAAATCG ATAAGCTTCC
ACCATG 56

(2) INFORMATION FOR SEQ ID NO:9:

10 (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 54 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

15 (ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

GATCAGAAGT TCCTATACTT TCTAGAGAAT AGGAACTTCG GAATAGGAAC TTCT
54

(2) INFORMATION FOR SEQ ID NO:10:

20 (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 41 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

25 (ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

GATCAGTTCC TATACTTTCT AGAGAATAGG AACTTCGGAA T
41

-52-

CLAIMS:

1. A DNA construct for transforming a eukaryotic cell, said construct comprising
a multifunctional DNA sequence flanked by two
5 directly repeated site-specific recombination sequences,
said multifunctional DNA sequence comprising a
gene encoding a selectable marker, and a DNA sequence of
interest,
wherein said DNA sequence of interest is flanked
10 by nucleotide sequences sharing homology to a nucleotide
sequence present in the eukaryotic cell, and the selectable
marker gene is operably linked to regulatory sequences
capable of expressing the gene in the eukaryotic cell.
2. The DNA construct of claim 1, wherein the
15 selectable marker gene and the DNA sequence of interest are
both flanked by nucleotide sequences sharing homology to a
nucleotide sequence present in the eukaryotic cell.
3. The DNA construct of claim 2, wherein the
multifunctional DNA sequence further comprises
20 a gene encoding a site specific recombinase
capable of recognizing the site-specific recombination
sequences flanking the multifunctional DNA sequence.
4. A DNA construct for inserting a DNA sequence of
interest into the DNA of a eukaryotic cell, said construct
25 comprising
a multifunctional DNA sequence flanked by direct
repeats of a first site-specific recombination sequence,
said multifunctional DNA sequence comprising a
gene encoding a first site specific recombinase capable of
30 recognizing the first site-specific recombination sequence,
and a DNA sequence targeted for insertion via homologous
recombination into the DNA of the eukaryotic cell,
said targeted DNA sequence being flanked by
nucleotide sequences sharing homology to nucleotide
35 sequences present in the eukaryotic cell, and comprising

-53-

said DNA sequence of interest and a excisable selection region,

said excisable selection region being flanked by direct repeats of a second site-specific recombination sequence and comprising a gene encoding a selectable marker, and a gene encoding a second site specific recombinase gene capable of recognizing the second site-specific recombination sequence,

wherein each gene is linked to regulatory sequences capable of expressing the gene in the eukaryotic cell.

5. The DNA construct of claim 3 further comprising DNA sequences capable of replicating the DNA molecule in a bacterial host, and DNA sequences encoding a bacterial selectable marker.

6. The DNA construct of claim 4, wherein expression of at least one of the first or second site specific recombinase gene is regulated by an inducible promoter.

7. The DNA construct of claim 6, wherein the expression of the recombinase gene is regulated by the glucocorticoid system.

8. A method of targeting a DNA sequence of interest into a specific site of a host cell's DNA, said method comprising the steps of

introducing the DNA construct described in claim 1 into individual eukaryotic cells,

selecting for cells having said DNA construct integrated into the DNA of the eukaryotic cell,

eliminating any randomly inserted DNA constructs,

identifying and culturing cells having said DNA sequence of interest integrated into the cell's DNA via a homologous recombination event.

9. A method for the production of fertile, transgenic plants wherein the transgenic plant has a DNA

-54-

sequence of interest integrated at a predetermined DNA sequence of the plant, said method comprising the steps of introducing into plant cells a DNA construct comprising

5 a multifunctional DNA sequence flanked by two directly repeated site-specific recombination sequences, said multifunctional DNA sequence comprising a gene encoding a selectable marker, and a DNA sequence of interest,

10 wherein said DNA sequence of interest is flanked by nucleotide sequences sharing homology to the predetermined nucleotide sequence present in the plant cell, and the selectable marker gene is operably linked to regulatory sequences capable of expressing the gene in the plant cell,

15 selecting for plant cells having said DNA construct integrated into the DNA of the plant cell, eliminating randomly inserted DNA constructs through expression of a recombinase gene capable of initiating recombination at the site-specific recombinase sequences in the plant cells,

20 identifying cells having said DNA sequence of interest integrated into the plant's DNA via a homologous recombination event, and

25 culturing said identified cells to generate an entire plant.

10. The method of claim 9, wherein the plant cells are haploid.

30 11. A plant entity consisting essentially of a plant cell, seed or plant produced from the *in vitro* introduction of an exogenous DNA fragment into a plant cell by the method of claim 9.

35 12. A method of directly selecting for insertion of a DNA sequence of interest into a specific sequence of an organism's DNA said method comprising the steps of

introducing the DNA construct of claim 4 into the cells of said organism,

applying selection means to isolate those cells having said DNA construct integrated into the DNA of the cell,

removing any randomly inserted DNA constructs, applying selection means to isolate those cells having the targeted DNA sequence integrated into the organism's DNA via a homologous recombination event, removing the excisable selection region, and culturing the resultant cells to regenerate an entire organism.

13. The method of claim 12, wherein the randomly integrated DNA sequences are eliminated by inducing the inducible promoter of the first recombinase gene, and the excisable selection region is removed by inducing the inducible promoter of the second site specific recombinase gene.

14. The DNA construct of claim 1, wherein said targeted DNA sequence is flanked by sequences homologous to the alcohol dehydrogenase gene.

15. A DNA construct for transforming eukaryotic cells, said construct comprising a multifunctional DNA sequence flanked by two directly repeated site-specific recombination sequences, said multifunctional DNA sequence comprising a gene encoding a selectable marker, and a DNA sequence targeted for insertion via homologous recombination into the DNA of the eukaryotic cell, wherein the selectable marker gene is operably linked to regulatory sequences capable of expressing the gene in the eukaryotic cell, and wherein

said targeted DNA sequence itself is flanked by nucleotide sequences sharing homology to nucleotide

-56-

sequences present in the eukaryotic cell, and comprises at least one polylinker region.

16. The recombinant DNA molecule of claim 15, wherein said multifunctional DNA sequence also comprises a gene
5 encoding a site specific recombinase capable of recognizing the site-specific recombination sequences flanking the multifunctional DNA sequence, the expression of which is regulated by an inducible promoter, and

- said DNA construct further comprises DNA
10 sequences capable of replicating the DNA molecule in a bacterial host, and DNA sequences encoding a bacterial selectable marker.

17. A kit comprising the construct of claim 16, and an inducer compound capable of inducing the inducible
15 promoter.

18. The method of claim 9, wherein the step of eliminating the randomly inserted DNA constructs is accomplished by including in the DNA construct a gene encoding a site specific recombinase capable of initiating
20 recombination at the site-specific recombination sequences flanking the multifunctional DNA sequence, said recombinase gene operably linked to sequences capable of expressing the recombinase gene in the plant cell.

19. The method of claim 9, wherein the step of
25 eliminating the randomly inserted DNA constructs is accomplished by

- including in the DNA construct a gene encoding a site specific recombinase capable of initiating recombination at the site-specific recombination sequences
30 flanking the multifunctional region, said recombinase gene operably linked to an inducible promoter, and

inducing the expression of the recombinase gene.

20. A method for the production of fertile, transgenic plants wherein the transgenic plant has a DNA

-57-

sequence of interest integrated at a predetermined DNA sequence of the plant, said method comprising the steps of introducing into plant cells a DNA construct comprising

- 5 a multifunctional DNA sequence flanked by two directly repeated site-specific recombination sequences, said multifunctional DNA sequence comprising a gene encoding a selectable marker, and a DNA sequence of interest, wherein the region of DNA encoding the selectable
- 10 marker and the DNA sequence of interest is flanked by nucleotide sequences sharing homology to the predetermined nucleotide sequence present in the plant cell, and the selectable marker gene is operably linked to regulatory sequences capable of expressing the gene in the plant cell,
- 15 eliminating randomly inserted DNA constructs through expression of a recombinase gene capable of initiating recombination at the site-specific recombinase sequences in the plant cells, selecting for plant cells having said DNA
- 20 sequence of interest integrated into the plant's DNA via a homologous recombination event, and culturing said identified cells to generate an entire plant.

21. The method of claim 20, wherein the step of
- 25 eliminating the randomly inserted DNA constructs is accomplished by including in the DNA construct a gene encoding a site specific recombinase capable of initiating recombination at the site-specific recombination sequences flanking the multifunctional region, said
- 30 recombinase gene operably linked to sequences capable of expressing the recombinase gene in the plant cell.

22. A plant entity consisting essentially of a plant cell, seed or plant produced from the *in vitro* introduction of an exogenous DNA fragment into a plant cell by the
- 35 method of claim 20.

AMENDED CLAIMS

[received by the International Bureau
on 8 July 1994 (08.07.94); original claims 1,4, 9,15 and 20 amended;
remaining claims unchanged (6 pages)]

1. A DNA construct for transforming a eukaryotic cell, said construct comprising
a multifunctional DNA sequence flanked by a pair
5 of directly repeated site-specific recombination sequences, said multifunctional DNA sequence comprising a gene encoding a selectable marker and a DNA sequence of interest, wherein said DNA sequence of interest is flanked by nucleotide sequences sharing homology to a nucleotide
10 sequence present in the eukaryotic cell, and the selectable marker gene is operably linked to regulatory sequences capable of expressing the gene in the eukaryotic cell.
2. The DNA construct of claim 1, wherein the selectable marker gene and the DNA sequence of interest are
15 both flanked by nucleotide sequences sharing homology to a nucleotide sequence present in the eukaryotic cell.
3. The DNA construct of claim 2, wherein the multifunctional DNA sequence further comprises a gene encoding a site specific recombinase capable of recognizing
20 the site-specific recombination sequences flanking the multifunctional DNA sequence.
4. A DNA construct for inserting a DNA sequence of interest into the DNA of a eukaryotic cell, said construct comprising a multifunctional DNA sequence flanked by a pair
25 of directly repeated first site-specific recombination sequences,
said multifunctional DNA sequence comprising a gene encoding a first site specific recombinase capable of recognizing the first site-specific recombination
30 sequences, and a DNA sequence targeted for insertion via homologous recombination into the DNA of the eukaryotic cell,
said targeted DNA sequence being flanked by
nucleotide sequences sharing homology to nucleotide
35 sequences present in the eukaryotic cell, and comprising

-59-

said DNA sequence of interest and an excisable selection region,

5 said excisable selection region being flanked by a pair of directly repeated second site-specific recombination sequences and comprising a gene encoding a selectable marker, and a gene encoding a second site specific recombinase gene capable of recognizing the second site-specific recombination sequences, wherein each gene is linked to regulatory sequences capable of expressing the
10 gene in the eukaryotic cell.

5. The DNA construct of claim 3 further comprising DNA sequences capable of replicating the DNA molecule in a bacterial host, and DNA sequences encoding a bacterial selectable marker.

15 6. The DNA construct of claim 4, wherein expression of at least one of the first or second site specific recombinase gene is regulated by an inducible promoter.

7. The DNA construct of claim 6, wherein the expression of the recombinase gene is regulated by the
20 glucocorticoid system.

8. A method of targeting a DNA sequence of interest into a specific site of a host cell's DNA, said method comprising the steps of

25 introducing the DNA construct described in claim 1 into individual eukaryotic cells, selecting for cells having said DNA construct integrated into the DNA of the eukaryotic cell, eliminating any randomly inserted DNA constructs, identifying and culturing cells having said DNA
30 sequence of interest integrated into the cell's DNA via a homologous recombination event.

9. A method for the production of fertile, transgenic plants wherein the transgenic plant has a DNA sequence of interest integrated at a predetermined DNA
35 sequence of the plant, said method comprising the steps of

-60-

introducing into plant cells a DNA construct comprising a multifunctional DNA sequence flanked by a pair of directly repeated site-specific recombination sequences, said multifunctional DNA sequence comprising a gene
5 encoding a selectable marker, and a DNA sequence of interest, wherein said DNA sequence of interest is flanked by nucleotide sequences sharing homology to the predetermined nucleotide sequence present in the plant cell, and the selectable marker gene is operably linked to
10 regulatory sequences capable of expressing the gene in the plant cell,

selecting for plant cells having said DNA construct integrated into the DNA of the plant cell, eliminating randomly inserted DNA constructs
15 through expression of a recombinase gene capable of initiating recombination at the site-specific recombinase sequences in the plant cells,

identifying cells having said DNA sequence of interest integrated into the plant's DNA via a homologous
20 recombination event, and

culturing said identified cells to generate an entire plant.

10. The method of claim 9, wherein the plant cells are haploid.

25 11. A plant entity consisting essentially of a plant cell, seed or plant produced from the *in vitro* introduction of an exogenous DNA fragment into a plant cell by the method of claim 9.

30 12. A method of directly selecting for insertion of a DNA sequence of interest into a specific sequence of an organism's DNA said method comprising the steps of

introducing the DNA construct of claim 4 into the cells of said organism,

applying selection means to isolate those cells having said DNA construct integrated into the DNA of the cell,

- 5 removing any randomly inserted DNA constructs, applying selection means to isolate those cells having the targeted DNA sequence integrated into the organism's DNA via a homologous recombination event, removing the excisable selection region, and culturing the resultant cells to regenerate an
10 entire organism.

13. The method of claim 12, wherein the randomly integrated DNA sequences are eliminated by inducing the inducible promoter of the first recombinase gene, and the excisable selection region is removed by inducing the
15 inducible promoter of the second site specific recombinase gene.

14. The DNA construct of claim 1, wherein said targeted DNA sequence is flanked by sequences homologous to the alcohol dehydrogenase gene.

- 20 15. A DNA construct for transforming eukaryotic cells, said construct comprising a multifunctional DNA sequence flanked by a pair of directly repeated site-specific recombination sequences, said multifunctional DNA sequence comprising a gene encoding a selectable marker,
25 and a DNA sequence targeted for insertion via homologous recombination into the DNA of the eukaryotic cell, wherein the selectable marker gene is operably linked to regulatory sequences capable of expressing the gene in the eukaryotic cell, and wherein said targeted DNA sequence itself is
30 flanked by nucleotide sequences sharing homology to nucleotide sequences present in the eukaryotic cell, and comprises at least one polylinker region.

16. The recombinant DNA molecule of claim 15, wherein said multifunctional DNA sequence also comprises a gene
35 encoding a site specific recombinase capable of recognizing

-62-

the site-specific recombination sequences flanking the multifunctional DNA sequence, the expression of which is regulated by an inducible promoter, and said DNA construct further comprises DNA sequences capable of replicating the DNA molecule in a bacterial host, and DNA sequences encoding a bacterial selectable marker.

17. A kit comprising the construct of claim 16, and an inducer compound capable of inducing the inducible promoter.

18. The method of claim 9, wherein the step of eliminating the randomly inserted DNA constructs is accomplished by including in the DNA construct a gene encoding a site specific recombinase capable of initiating recombination at the site-specific recombination sequences flanking the multifunctional DNA sequence, said recombinase gene operably linked to sequences capable of expressing the recombinase gene in the plant cell.

19. The method of claim 9, wherein the step of eliminating the randomly inserted DNA constructs is accomplished by

including in the DNA construct a gene encoding a site specific recombinase capable of initiating recombination at the site-specific recombination sequences flanking the multifunctional region, said recombinase gene operably linked to an inducible promoter, and inducing the expression of the recombinase gene.

20. A method for the production of fertile, transgenic plants wherein the transgenic plant has a DNA sequence of interest integrated at a predetermined DNA sequence of the plant, said method comprising the steps of introducing into plant cells a DNA construct comprising a multifunctional DNA sequence flanked by a pair of directly repeated site-specific recombination sequences, said multifunctional DNA sequence comprising a gene encoding a selectable marker, and a DNA sequence of

-63-

- interest, wherein the region of DNA encoding the selectable marker and the DNA sequence of interest is flanked by nucleotide sequences sharing homology to the predetermined nucleotide sequence present in the plant cell, and the
- 5 selectable marker gene is operably linked to regulatory sequences capable of expressing the gene in the plant cell, eliminating randomly inserted DNA constructs through expression of a recombinase gene capable of initiating recombination at the site-specific recombinase
- 10 sequences in the plant cells, selecting for plant cells having said DNA sequence of interest integrated into the plant's DNA via a homologous recombination event, and culturing said identified cells to generate an
- 15 entire plant.
21. The method of claim 20, wherein the step of eliminating the randomly inserted DNA constructs is accomplished by including in the DNA construct a gene encoding a site specific recombinase capable of initiating
- 20 recombination at the site-specific recombination sequences flanking the multifunctional region, said recombinase gene operably linked to sequences capable of expressing the recombinase gene in the plant cell.
22. A plant entity consisting essentially of a plant
- 25 cell, seed or plant produced from the *in vitro* introduction of an exogenous DNA fragment into a plant cell by the method of claim 20.

-64-

STATEMENT UNDER ARTICLE 19

Prosecution of the priority application has led to the amendment of several claims, and these amendments are reflected in the substitute sheets. Claims 1, 4, 9, 15 and 20 are amended to specify that the multifunctional DNA sequence is flanked by a pair of directly repeated site-specific recombination sequences. This amendment is made to remove an ambiguity regarding the number and location of the site-specific recombination sequences. During prosecution of the priority application, the Examiner requested such an amendment to clarify that a site-specific recombination sequence is located at the start of the multifunctional DNA sequence and a second directly repeated site-specific recombination sequence is located at the other end of the multifunctional DNA sequence region. Applicants submit herewith replacement sheets for pages 52-57 of the International Application as originally filed.

1 / 11




FIG. 1A
SUBSTITUTE SHEET (RULE 26)

2 / 11




FIG. 1B

SUBSTITUTE SHEET (RULE 26)







FIG.3A

SUBSTITUTE SHEET (RULE 26)



5 / 11

FIG.3B



SUBSTITUTE SHEET (RULE 26)

FIG. 4A

mRNA exon BglIII AGATCT intron GUS

pUbiGUS

GATCAGAAGTTCCTATACTTTCTAGAGAATAGGAACTTCGGAATAGGAATTCT

XbaI

*** **

GATCAGTTCCATACTTTCTAGAGAATAGGAACTTCGGAAT

*** **

XbaI


pUFRTG

pUFRTmg

The diagram illustrates the GUS reporter gene construct and its excision products. The top part shows the full construct, which includes a Ubiquitin promoter, exons, introns, FRT sites, a 1.3kb pNEOβGAL fragment, and a GUS coding sequence with a poly A site. The bottom part shows two excision products: one with a functional GUS coding sequence (pU2FRTG) and one with a defective GUS coding sequence (pU2FRTGrev).


7/11

FIG. 5



SUBSTITUTE SHEET (RULE 26)

FIG. 6



SUBSTITUTE SHEET (RULE 26)

FIG.7






FIG. 8





INTERNATIONAL SEARCH REPORT

International application no.
PCT/US94/00927

A. CLASSIFICATION OF SUBJECT MATTER

IPC(5) : C12N 5/00, 15/00; A01H 1/04, 5/00, 5/10
US CL : 435/172.1, 172.3, 240.4; 800/200, 205, 250; 935/18

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/172.1, 172.3, 240.4; 800/200, 205, 250; 935/18

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
APS, BIOSIS, CA

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	MOL GEN GENET, Volume 223, issued 1990, J. Odell et al, "Site-directed recombination in the genome of transgenic tobacco", pages 369-378, especially pages 369, 372, 376, 377.	1-22
Y	PROC. NATL. ACAD. SCI., USA, Volume 88, issued December 1991, E. C. Dale et al, "Gene transfer with subsequent removal of the selection gene from the host genome", pages 10558-10562, especially page 10558, Abstract.	1-22
Y	SCIENCE, Volume 244, issued 1989, M. R. Capecchi, "Altering the Genome by Homologous Recombination", pages 1288-1292, especially pages 1288 (Abstract) and 1292.	1-8, 12-19

☒ Further documents are listed in the continuation of Box C.
 ☐ See patent family annex.

* "A" document defining the general state of the art which is not considered to be part of particular relevance	* "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
* "E" earlier document published on or after the international filing date	* "X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
* "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	* "Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
* "O" document referring to an oral disclosure, use, exhibition or other means	* "Z" document member of the same patent family
* "P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

29 APRIL 1994

Date of mailing of the international search report

MAY 11 1994

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703)305-3230

Authorized officer

Erich E. Veitenheimer

Telephone No. (703)308-0196

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US94/00927

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	NUCLEIC ACIDS RESEARCH, Volume 14, Number 5, issued 1986, R. H. Hoess et al, "The role of the loxP spacer region in P1 site-specific recombination", pages 2287-2300, especially pages 2292 and 2299.	1-8, 12-19
Y	THE PLANT CELL, Volume 2, issued July 1990, W. J. Gordon-Kamm et al, "Transformation of Maize Cells and Regeneration of Fertile Transgenic Plants", pages 603-618, see entire article.	9-11, 20-22